

**q -IDENTITIES AND AFFINIZED PROJECTIVE VARIETIES
II. FLAG VARIETIES**

PETER BOUWKNEGT AND NICK HALMAGYI

Dedicated to the memory of Prof. H.S. Green

ABSTRACT. In a previous paper we defined the concept of an affinized projective variety and its associated Hilbert series. We computed the Hilbert series for varieties associated to quadratic monomial ideals. In this paper we show how to apply these results to affinized flag varieties. We discuss various examples and conjecture a correspondence between the Hilbert series of an affinized flag variety and a modified Hall-Littlewood polynomial. We briefly discuss the application of these results to quasi-particle character formulas for affine Lie algebra modules.

1. INTRODUCTION

In a previous paper [Bo], one of us introduced the concept of an affinized projective variety and the associated Hilbert series of its coordinate ring. It was shown how q -identities naturally arise from different ways of computing this Hilbert series. This was made explicit in the case of projective varieties defined by quadratic monomial ideals, where, on the one hand, an explicit basis for the coordinate ring of the associated affinized projective variety led to an expression of the Hilbert series, while, on the other hand, an algorithm was described how to obtain an alternating sum formula (resembling an ‘affinization’ of Taylor’s resolution for monomial ideals) for the same Hilbert series.

In the present paper we will show how one can apply the results of [Bo] to compute the Hilbert series of an affinized flag variety. The ideal that defines the coordinate ring of a flag variety, albeit quadratic, is not a monomial ideal. We will show, however, how one can in principle construct a projective variety defined by a quadratic monomial ideal which has the *same* Hilbert series. This involves two ideas. The first is to consider the (monomial) ideal of leading terms $\langle \text{LT}(I) \rangle$ instead of I . The second idea is to remove the non-quadratic monomials by the addition of extra variables. We carry out this program in several examples.

The Hilbert series of a flag variety based on a group G , as well as the Hilbert series of its affinization, are naturally characters of $\text{Lie } G = \mathfrak{g}$. In fact, upto a trivial factor, the Hilbert

series of the affinized flag variety can be interpreted as a q -deformation of the character of a tensor product of $\mathfrak{g}$ -modules. One of the main results of this paper is a conjectured correspondence between the Hilbert series of an affinized flag variety and another such q -deformation, namely the modified Hall-Littlewood polynomial (or Milne polynomial). Modified Hall-Littlewood polynomials are related in various ways to characters of affine Lie algebras. Our results thus provide new, explicit, expressions for these characters. In fact, this was the main motivation for this work. A relation between the characters of affine Lie algebras and the geometry of infinite dimensional flag varieties was also pointed out and explored in [FS1,FS2].

The affine Lie algebra characters are naturally obtained in what is known as the ‘Universal Chiral Partition Function’ (UCPF) form, which is conjectured to be a universal form for the (chiral) characters of any two dimensional conformal field theory [BM], and is closely related to a description of that conformal field theory in terms of quasi-particles. The results of this paper further support the validity of this conjecture, at least in the case of conformal field theories based on affine Lie algebras (WZW models). In a separate paper [BCR] we will explore the results of this paper in connection with the exclusion statistics (cf. [Ha,BS4,GS] and reference therein) satisfied by these quasi-particles. The example of $\mathfrak{so}_5$, treated in detail in this paper, is particularly relevant with regards to possible applications to the quasi-particles (‘non-abelian electrons’) in $SO(5)$ superspin regimes of strongly correlated electrons on a two-leg ladder [BS3].

This paper is organized as follows. In section 2 we recall the definition of the homogeneous coordinate ring of a (finite dimensional) flag variety and show how one can, in principle, compute the ideal of quadratic relations. We illustrate this in a few examples. In section 3 we introduce affinized flag varieties and outline a procedure to compute the Hilbert series of such a variety. In section 4 we apply the procedure in a few examples, namely, $\mathfrak{sl}_n$, $n = 2, 3, 4$, and $\mathfrak{so}_{2n+1}$, $n = 2, 3$, and in section 5 we comment on the relation between the Hilbert series of an affinized flag variety and modified Hall-Littlewood polynomials as well as characters of integrable highest weight modules of affine Lie algebras. We also comment on the application of our results to theories of quasi-particles, their exclusion statistics and the UCPF.

Throughout this paper we will use the notation of [Bo].

2. FLAG VARIETIES

2.1. Generalities.

Most of the material in this section is quite standard. We refer to [FH,Fu] for detailed expositions.

Let $\mathfrak{g}$ be a (complex) finite dimensional simple Lie algebra of rank ℓ . Let $\{\alpha_i\}_{i=1}^\ell$ and $\{\Lambda_i\}_{i=1}^\ell$ denote the simple roots and fundamental weights of $\mathfrak{g}$, respectively. Let $L(\Lambda_i)$, $i = 1, \dots, \ell$, denote the finite dimensional irreducible representation of $\mathfrak{g}$ with highest weight Λ_i and dimension $D_i = \dim L(\Lambda_i)$. For any $\mathfrak{g}$ -module V , we denote by $\text{Sym}^M V$

the symmetrized tensor product of M copies of V , and by $\text{Sym}^\bullet V = \bigoplus_{M \geq 0} \text{Sym}^M V$ the symmetric algebra on V . The module

$$(2.1) \quad \text{Sym}^{M_1} L(\Lambda_1) \otimes \dots \otimes \text{Sym}^{M_\ell} L(\Lambda_\ell),$$

is completely reducible, for any choice of $M_i \geq 0$, and contains $L(M_1 \Lambda_1 + \dots + M_\ell \Lambda_\ell)$ as a submodule. Upon introducing coordinates $x_a^{(i)}$, $a = 1, \dots, D_i$, with respect to an orthonormal basis $v_a^{(i)}$ for each $L(\Lambda_i)$, $i = 1, \dots, \ell$, we can identify

$$(2.2) \quad \begin{aligned} \mathbf{S} &\cong \text{Sym}^\bullet L(\Lambda_1) \otimes \dots \otimes \text{Sym}^\bullet L(\Lambda_\ell), \\ \mathbf{S}_{(M_1, \dots, M_\ell)} &\cong \text{Sym}^{M_1} L(\Lambda_1) \otimes \dots \otimes \text{Sym}^{M_\ell} L(\Lambda_\ell), \end{aligned}$$

where $\mathbf{S} \equiv \mathbb{C}[x_a^{(i)}]_{a=1, \dots, D_i}^{i=1, \dots, \ell}$ and $\mathbf{S}_{(M_1, \dots, M_\ell)}$ denotes the subspace of polynomials in $\mathbf{S}$, homogeneous of degree M_i in the $\{x_a^{(i)}\}_{a=1, \dots, D_i}$. The Lie algebra $\mathfrak{g}$ is realized on $\mathbf{S}_{(M_1, \dots, M_\ell)}$ in terms of linear differential operators. Note, furthermore,

$$(2.3) \quad \dim \mathbf{S}_{(M_1, \dots, M_\ell)} = \prod_{i=1}^{\ell} \binom{M_i + D_i - 1}{D_i - 1}.$$

Of course, the main problem is to give a concrete description for how the irreducible module $L(M_1 \Lambda_1 + \dots + M_\ell \Lambda_\ell)$ is contained in $\mathbf{S}_{(M_1, \dots, M_\ell)}$. The solution is remarkably simple. By a theorem of Kostant (see, e.g., [LT]) we have that $\mathbf{S}$, as a $\mathfrak{g}$ -module, contains an ideal I generated by quadratic relations such that

$$(2.4) \quad L(M_1 \Lambda_1 + \dots + M_\ell \Lambda_\ell) \cong \mathbf{S}(V)_{(M_1, \dots, M_\ell)},$$

where $\mathbf{S}(V) = \mathbf{S}/I$. Furthermore, it turns out that $\mathbf{S}(V)$ is precisely the homogeneous coordinate ring of the (complete) flag variety of $\mathfrak{g}$, i.e., the variety $V = \mathbf{V}(I)$, corresponding to $I \subset \mathbb{C}[x_a^{(i)}]$, is isomorphic to G/B where G is the (complex) Lie group such that $\text{Lie } G = \mathfrak{g}$ and B is a Borel subgroup of G . More generally, omitting some of the variables $x_a^{(i)}$ for certain i (or, equivalently, taking $M_i = 0$ for some i), corresponds to the homogeneous coordinate ring of a partial flag variety G/P where P is a parabolic subgroup of G .

The ideal of quadratic relations I can be determined, in principle, by analyzing the tensor products $L(\Lambda_i) \otimes L(\Lambda_j)$ (see the examples in sections 2.2 and 2.3). It appears that explicit results are known only for $\mathfrak{g} = \mathfrak{sl}_n$ (or $\mathfrak{gl}_n$) and some low rank cases.

Besides the grading by the multi-degree $\mathbf{M} = (M_1, \dots, M_\ell)$, the $\mathbf{S}$ -module $\mathbf{S}(V)$ is also graded by the $\mathfrak{g}$ -weight. Denote by $\mathbf{S}(V)_{(\mathbf{M}; \lambda)}$ the space of homogeneous polynomials in $\mathbf{S}(V)$ of multi-degree $\mathbf{M}$ and $\mathfrak{g}$ -weight λ . The Hilbert function of $\mathbf{S}(V)$ is then defined as the (formal) $\mathfrak{g}$ -character

$$(2.5) \quad h_V(\mathbf{M}) = \sum_{\lambda} \dim \mathbf{S}(V)_{(\mathbf{M}; \lambda)} e^{\lambda},$$

or, by (2.4), as the character of the irreducible module $L(M_1\Lambda_1 + \dots + M_\ell\Lambda_\ell)$.

As discussed in [Bo], the Hilbert function $h_V(\mathbf{M})$ can be computed either by constructing an explicit basis for the module $\mathbf{S}(V)$ or by applying the Euler-Poincaré principle to a free resolution of $\mathbf{S}(V)$

$$(2.6) \quad 0 \rightarrow F^{(\nu)} \xrightarrow{d_\nu} \dots \xrightarrow{d_3} F^{(2)} \xrightarrow{d_2} F^{(1)} \xrightarrow{d_1} F^{(0)} \cong \mathbf{S} \rightarrow \mathbf{S}(V) \rightarrow 0,$$

where the maps d_i intertwine with the action of $\mathfrak{g}$. Here, $F^{(j)} = \bigoplus_k \mathbf{S}(-\mathbf{a}_k^{(j)}; -\mu_k^{(j)})$ for some set of vectors $\mathbf{a}_k^{(j)} = ((a_k^{(j)})_1, \dots, (a_k^{(j)})_\ell)$ with positive integer coefficients, and integral weights $\mu_k^{(j)}$. In fact, because of the $\mathfrak{g}$ structure, the weights $\mu_k^{(j)}$ nicely combine into weights of $\mathfrak{g}$ -modules, i.e., we can write $F^{(j)} = \bigoplus_k (\mathbf{S}(-\mathbf{a}_k^{(j)}) \otimes V_k^{(j)})$ for some set of (finite dimensional) $\mathfrak{g}$ -modules $V_k^{(j)}$. Furthermore, as in [Bo], for any module M we have denoted by $M(\mathbf{a}; \mu)$ the same module with multi-degree and weight shifted by $\mathbf{a}$ and μ , respectively. Application of the Euler-Poincaré principle to (2.6) yields the following alternating sum formula for the character $h_V(M_1, \dots, M_\ell)$ of the irreducible module $L(M_1\Lambda_1 + \dots + M_\ell\Lambda_\ell)$

$$(2.7) \quad h_V(M_1, \dots, M_\ell) = \sum_j (-1)^j \sum_k h_{\mathbf{S}}(M_1 - (a_k^{(j)})_1, \dots, M_\ell - (a_k^{(j)})_\ell) \chi_{V_k^{(j)}},$$

where χ_V denotes the (formal) character of the $\mathfrak{g}$ -module V and

$$(2.8) \quad h_{\mathbf{S}}(M_1, \dots, M_\ell) = \sum_{\lambda} \dim \mathbf{S}_{(M_1, \dots, M_\ell; \lambda)} e^\lambda = \sum_{\sum_a m_a^{(i)} = M_i} e^{\sum_{i,a} m_a^{(i)} \lambda_a^{(i)}},$$

denotes the character valued Hilbert function of $\mathbf{S} = \mathbb{C}[x_a^{(i)}]$. The weights of $L(\Lambda_i)$ are denoted by $\lambda_a^{(i)}$, $a = 1, \dots, D_i$.

The existence of a (finite) free resolution (2.6) is guaranteed by Hilbert's syzygy theorem, but explicit descriptions are known apparently only in special examples. Some examples will be discussed in sections 2.2 and 2.3. Based on these and other examples we have

Conjecture 2.1. *Consider the minimal resolution (2.6) of $\mathbf{S}(V)$. Let ν be the length of this resolution, $D = \dim G/B$, $\ell = \text{rank } \mathfrak{g}$ and $D_i = \dim L(\Lambda_i)$, then*

- i. $\sum_{i=1}^{\ell} D_i = D + \ell + \nu$
- ii. *The minimal resolution is symmetric under simultaneous interchange of $\mathbf{S}(a_1, \dots, a_\ell; \mu) \leftrightarrow \mathbf{S}(2 - D_1 - a_1, \dots, 2 - D_\ell - a_\ell; -\mu)$ (or, equivalently, $\mathbf{S}(a_1, \dots, a_\ell) \otimes V \leftrightarrow \mathbf{S}(2 - D_1 - a_1, \dots, 2 - D_\ell - a_\ell) \otimes V^*$, where V^* is the $\mathfrak{g}$ -module contragredient to V), and reversal of all arrows.*

In particular, since $F^{(0)} = \mathbf{S} = \mathbf{S}(0, \dots, 0; 0)$ it follows from ii. that $F^{(\nu)} \cong \mathbf{S}(2 - D_1, \dots, 2 - D_\ell; 0)$.

In the remainder of this section we will explicitly go through some examples in preparation for the affinization of this construction which will be discussed in sections 3 and 4.

2.2. Example: $\mathfrak{sl}_n$.

Consider the Lie algebra $A_{n-1} \cong \mathfrak{sl}_n$ of rank $\ell = n - 1$. In terms of an overcomplete basis $\{\epsilon_i\}_{i=1}^n$ of $\mathbb{C}^{n-1}$, satisfying

$$(2.9) \quad (\epsilon_i, \epsilon_j) = \delta_{ij} - \frac{1}{n}, \quad \sum_{i=1}^n \epsilon_i = 0,$$

the simple roots and fundamental weights of $\mathfrak{sl}_n$ can be written as

$$(2.10) \quad \alpha_i = \epsilon_i - \epsilon_{i+1}, \quad \Lambda_i = \epsilon_1 + \dots + \epsilon_i, \quad i = 1, \dots, n-1.$$

The irreducible finite dimensional representation $L(\Lambda_i)$, of dimension $D_i = \binom{n}{i}$, has weights

$$(2.11) \quad \epsilon_{i_1} + \dots + \epsilon_{i_k}, \quad 1 \leq i_1 < \dots < i_k \leq n.$$

Let us denote the coordinate corresponding to the vector of weight $\epsilon_{i_1} + \dots + \epsilon_{i_k}$ by $x_{i_1 \dots i_k}$. For convenience we extend the definition of $x_{i_1 \dots i_k}$ to arbitrary sequences $1 \leq i_1, \dots, i_k \leq n$ by anti-symmetry, and raise and lower indices by means of the ϵ -tensor, i.e.,

$$(2.12) \quad x^{i_1 \dots i_{n-k}} = \frac{1}{k!} \epsilon^{i_1 \dots i_{n-k} j_1 \dots j_k} x_{j_1 \dots j_k}.$$

We have a realization of $\mathfrak{gl}_n$ on $\mathbf{S} = \mathbb{C}[x_{i_1}, \dots, x_{i_1 \dots i_{n-1}}]$ in terms of linear differential operators, given by

$$(2.13) \quad e_{ij} = x_i \frac{\partial}{\partial x_j} + \sum_k x_{ik} \frac{\partial}{\partial x_{jk}} + \dots + \sum_{k_1 < \dots < k_{n-2}} x_{ik_1 \dots k_{n-2}} \frac{\partial}{\partial x_{jk_1 \dots k_{n-2}}}.$$

That is, the operators (2.13) satisfy the defining relations of $\mathfrak{gl}_n$

$$(2.14) \quad [e_{ij}, e_{kl}] = \delta_{jk} e_{il} - \delta_{il} e_{kj}.$$

Henceforth, we consider $\mathbf{S}$ as an $\mathfrak{sl}_n$ -module. As argued before, the $\mathfrak{sl}_n$ -module $\mathbf{S}_{(M_1, \dots, M_{n-1})}$ contains $L(\Lambda) \equiv L(M_1 \Lambda_1 + \dots + M_{n-1} \Lambda_{n-1})$ as a submodule. In particular, the highest weight vector of $L(\Lambda)$ is given by $x_1^{M_1} x_{12}^{M_2} \dots x_{12 \dots n-1}^{M_{n-1}}$.

It is well-known (see, e.g., [FH, Fu]) that the ideal I of quadratic relations is generated by the polynomials $\{\sigma_{i_2 \dots i_k}^{j_1 \dots j_{n-l-1}} \mid 1 \leq k \leq l < n\}$, where

$$(2.15) \quad \sigma_{i_2 \dots i_k}^{j_1 \dots j_{n-l-1}} = \epsilon^{i_1 k_1 \dots k_l j_1 \dots j_{n-l-1}} x_{i_1 i_2 \dots i_k} x_{k_1 k_2 \dots k_l},$$

Note that the invariant subspace of polynomials $\sigma_{i_2 \dots i_k}^{j_1 \dots j_{n-l-1}}$, for fixed $1 \leq k \leq l < n$, is isomorphic to $L(\Lambda_{k-1}) \otimes L(\Lambda_{l+1})$ and will thus, in general, be reducible under $\mathfrak{sl}_n$.

Remark 2.1. We briefly comment on the identification of the coordinate ring defined above with that of a flag variety in the case of $\mathfrak{sl}_n$ (see [Fu] for more details). Let $E = \mathbb{C}^n$. Fix a sequence $n \geq d_s > \dots > d_2 > d_1 \geq 0$. The (partial) flag variety $\mathrm{Fl}^{d_1 \dots d_s}(E)$ is the set of flags

$$(2.16) \quad \{E_s \subset \dots \subset E_2 \subset E_1 \subset E \mid \mathrm{codim} E_i = d_i, 1 \leq i \leq s\}.$$

The group $G = GL(E) = GL(n, \mathbb{C})$ acts transitively on $\mathrm{Fl}^{d_1 \dots d_s}(E)$. Now, let $\{e_i\}_{i=1}^n$ be a basis of E and let $F^{(0)} = \{E_s^{(0)} \subset \dots \subset E_2^{(0)} \subset E_1^{(0)} \subset E\}$ be the flag defined by $E_i^{(0)} = \langle e_{d_i+1}, e_{d_i+2}, \dots, e_n \rangle$. Furthermore, let P be the parabolic subgroup of $GL(E)$ fixing the flag $F^{(0)}$, i.e.,

$$(2.17) \quad P = \{g \in GL(E) \mid g(E_i) \subset E_i, 1 \leq i \leq s\},$$

then we can identify $\mathrm{Fl}^{d_1 \dots d_s}(E) = G/P$. Specifically, note that a matrix in P has invertible matrices in blocks of sizes $d_1, d_2 - d_1, \dots, d_s - d_{s-1}, n - d_s$ down the diagonal, with arbitrary entries below these blocks and vanishing entries above. For the complete flag variety $\mathrm{Fl}^{1,2,\dots,n}(E)$ we have $P = B$ where B is a Borel subgroup of G . We have a natural embedding, the so-called Plücker embedding,

$$(2.18) \quad \mathrm{Fl}^{d_1 \dots d_s}(E) \hookrightarrow \mathbb{P}(\bigwedge^{d_1} E) \times \dots \times \mathbb{P}(\bigwedge^{d_s} E).$$

Since $\bigwedge^p E$ has a basis $e_{i_1} \wedge \dots \wedge e_{i_p}$, we have a natural set of homogeneous coordinates $x_{i_1 \dots i_p}$ for $\mathbb{P}(\bigwedge^p E)$. The relations (2.15) are precisely the “Plücker relations” for the embedding (2.18).

For $\mathfrak{sl}_n$ the (minimal) resolution (2.6) of $\mathbf{S}(V) = \mathbf{S}/I$ does not seem to be known for general n . We discuss some examples for small n .

2.2.1. $\mathfrak{sl}_2$.

For $\mathfrak{sl}_2$ we have a single fundamental representation $L(\Lambda_1)$ of dimension $D_1 = 2$. Moreover, $\mathrm{Sym}^2 L(\Lambda_1) \cong L(2\Lambda_1)$, so $\mathbf{S} = \mathbb{C}[x_1, x_2]$ while the ideal I is trivial. We immediately conclude that the character of the irreducible (spin- $M/2$) module $L(M\Lambda_1)$ is given by (cf. (2.7))

$$(2.19) \quad h(M) = \sum_{m_1+m_2=M} e^{m_1\epsilon_1+m_2\epsilon_2} = \sum_{m_1+m_2=M} e^{(m_1-m_2)\epsilon_1},$$

as one may readily verify.

2.2.2. $\mathfrak{sl}_3$.

We have two fundamental representations $L(\Lambda_i)$, $i = 1, 2$, of dimension $D_1 = D_2 = 3$, hence $\mathbf{S} = \mathbb{C}[x_i, x^i]_{i=1,2,3}$. From

$$(2.20) \quad \begin{aligned} \mathrm{Sym}^2 L(\Lambda_1) &\cong L(2\Lambda_1), \\ L(\Lambda_1) \otimes L(\Lambda_2) &\cong L(\Lambda_1 + \Lambda_2) \oplus L(0), \\ \mathrm{Sym}^2 L(\Lambda_2) &\cong L(2\Lambda_2), \end{aligned}$$

it follows that I is generated by a single ($\mathfrak{sl}_3$ -singlet) generator σ . Explicitly,

$$(2.21) \quad \sigma = x_i x^i = \frac{1}{2} \epsilon^{ijk} x_i x_{jk}.$$

The resolution of $\mathbf{S}(V) = \mathbf{S}/I$ is obviously given by

$$(2.22) \quad 0 \rightarrow F^{(1)} \xrightarrow{d_1} F^{(0)} \rightarrow \mathbf{S}/I \rightarrow 0,$$

where $F^{(1)} \cong \mathbf{S}(-1, -1)$ and $d_1 : F^{(1)} \rightarrow F^{(0)}$ is defined by

$$(2.23) \quad d_1 : e^{(1)} \mapsto \sigma e^{(0)},$$

where $e^{(i)}$ is the generator of $F^{(i)}$. See also Fig. 2.1.

$$\begin{array}{ccc} F^{(0)} & & \mathbf{S}_{(0,0)} \\ \uparrow d_1 & & \uparrow \\ F^{(1)} & & \mathbf{S}_{(-1,-1)} \end{array}$$

Fig. 2.1. Resolution of $\mathbf{S}(V)$ for $\mathfrak{sl}_3$

The resolution (2.22) leads to the following well-known expression for the character $h_V(M_1, M_2)$ of the irreducible module $L(M_1\Lambda_1 + M_2\Lambda_2)$

$$(2.24) \quad h_V(M_1, M_2) = h_{\mathbf{S}}(M_1, M_2) - h_{\mathbf{S}}(M_1 - 1, M_2 - 1),$$

where

$$(2.25) \quad h_{\mathbf{S}}(M_1, M_2) = \sum_{\substack{m_1 + m_2 + m_3 = M_1 \\ m_1 + m_2 + m_3 = M_2}} e^{\sum_i (m_i - m^i) \epsilon_i},$$

denotes the Hilbert function of $\mathbf{S}$.

2.2.3. $\mathfrak{sl}_4$.

For $\mathfrak{sl}_4$ we have three fundamental representations of dimensions $D_1 = 4$, $D_2 = 6$ and $D_3 = 4$. The relations generating $I \subset \mathbb{C}[x_i, x_{ij}, x_{ijk}]$ arise from the tensor products

$$\begin{aligned}
 L(\Lambda_1) \otimes L(\Lambda_3) &\cong L(\Lambda_1 + \Lambda_3) \oplus L(0), \\
 \text{Sym}^2 L(\Lambda_2) &\cong L(2\Lambda_2) \oplus L(0), \\
 L(\Lambda_1) \otimes L(\Lambda_2) &\cong L(\Lambda_1 + \Lambda_2) \oplus L(\Lambda_3), \\
 L(\Lambda_2) \otimes L(\Lambda_3) &\cong L(\Lambda_2 + \Lambda_3) \oplus L(\Lambda_1),
 \end{aligned}
 \tag{2.26}$$

and are given by

$$\begin{aligned}
 \sigma &= \epsilon^{ijkl} x_i x_{jkl}, \\
 \bar{\sigma} &= \epsilon^{ijkl} x_{ij} x_{kl}, \\
 \sigma^i &= \epsilon^{ijkl} x_j x_{kl}, \\
 \sigma_i &= \epsilon^{jklm} x_{ijk} x_{lm}.
 \end{aligned}
 \tag{2.27}$$




Fig. 2.2. Resolution of $\mathbf{S}(V)$ for $\mathfrak{sl}_4$.

The resolution of $\mathbf{S}(V) = \mathbf{S}/I$ is depicted in Fig. 2.2. We refrain from giving the explicit maps which are easily worked out from the shifts in degrees indicated in the figure. Also, we have used the notation $\mathbf{S}(a, b)_V = \mathbf{S}(a, b) \otimes V$ for an $\mathfrak{sl}_4$ module V . Note that the resolution in is complete agreement with Conjecture 2.1.

2.3. Example: $\mathfrak{so}_{2n+1}$.

Consider the Lie algebra $B_n \cong \mathfrak{so}_{2n+1}$ of rank $\ell = n$. Let $\{\epsilon_i\}_{i=1}^n$ be an orthonormal basis of $\mathbb{C}^n$. The simple roots and fundamental weights of $\mathfrak{so}_{2n+1}$ can be written as

$$(2.28) \quad \begin{aligned} \alpha_i &= \epsilon_i - \epsilon_{i+1}, & \Lambda_i &= \epsilon_1 + \dots + \epsilon_i, & i &= 1, \dots, n-1, \\ \alpha_n &= \epsilon_n, & \Lambda_n &= \frac{1}{2}(\epsilon_1 + \dots + \epsilon_n). \end{aligned}$$

The irreducible finite dimensional representations $L(\Lambda_i)$ of $\mathfrak{so}_{2n+1}$ have dimension

$$(2.29) \quad \begin{aligned} D_i &= \binom{2n+1}{i}, & i &= 1, \dots, n-1, \\ D_n &= 2^n. \end{aligned}$$

For simplicity, we only consider here the representations $L(\Lambda_1)$ and $L(\Lambda_n)$, i.e., the vector and the spinor representation, respectively. The weights of $L(\Lambda_1)$ are

$$(2.30) \quad \{\epsilon_1, \epsilon_2, \dots, \epsilon_n, 0, -\epsilon_n, \dots, -\epsilon_1\},$$

with corresponding coordinates $\{x_i\} = \{x_1, \dots, x_n, x_0, x_{\bar{n}}, \dots, x_{\bar{1}}\}$, while the weights of $L(\Lambda_n)$ are

$$(2.31) \quad \left\{ \frac{1}{2}(\pm\epsilon_1 \pm \epsilon_2 \pm \dots \pm \epsilon_n) \right\},$$

with corresponding coordinates $\{x_\alpha\} = \{x_{\pm\pm\dots\pm}\}$. The explicit realization of $\mathfrak{so}_{2n+1}$ on $\text{Sym}^\bullet L(\lambda_1) \otimes \text{Sym}^\bullet L(\Lambda_n) \cong \mathbb{C}[x_i, x_\alpha]$ is easily constructed (see section 2.3.1 for $\mathfrak{so}_5$). The relevant tensor products are

$$(2.32) \quad \begin{aligned} \text{Sym}^2 L(\Lambda_1) &\cong L(2\Lambda_1) \oplus L(0), \\ L(\Lambda_1) \otimes L(\Lambda_n) &\cong L(\Lambda_1 + \Lambda_n) \oplus L(\Lambda_n), \\ \text{Sym}^2 L(\Lambda_n) &\cong L(2\Lambda_n) \oplus \left(\bigoplus_{\substack{k=1 \\ \text{restrictions}}}^{n-1} L(\Lambda_k) \right), \end{aligned}$$

where, in the last tensor product, the sum is over those k such that $k \equiv n \pmod{4}$ or $k \equiv (n+1) \pmod{4}$. The first two of these lead to quadratic relations of the form

$$(2.33) \quad \sigma = g^{ij} x_i x_j, \quad \sigma_\alpha = (\gamma^i)_\alpha{}^\beta x_i x_\beta,$$

for some matrix g^{ij} and some set of γ -matrices γ^i , while the last leads to relations depending on the specific value of n . The realization of $\mathfrak{so}_{2n+1}$ can be chosen in such a way that the matrices satisfy the properties

$$(2.34) \quad \{\gamma^i, \gamma^j\} = 2g^{ij} = \delta^{i\bar{j}}, \quad (\gamma^i)_\alpha{}^\beta = (\gamma^{\bar{i}})_\beta{}^\alpha.$$

In addition, it is possible to define a charge conjugation matrix $C^{\alpha\beta}$, so that if spinor indices are raised and lowered by means of $C^{\alpha\beta}$ and its inverse, e.g., $x^\alpha = C^{\alpha\beta} x_\beta$, then we have the additional symmetry property

$$(2.35) \quad (\gamma^i)^{\alpha\beta} = -(\gamma^i)^{\beta\alpha}.$$

In the next subsection we discuss the case $\mathfrak{so}_5$ in somewhat more detail.

2.3.1. $\mathfrak{so}_5$.

Consider the Lie algebra $\mathfrak{so}_5$. As discussed, we introduce variables $\{x_i\} = \{x_1, x_2, x_0, x_{\bar{2}}, x_{\bar{1}}\}$ and $\{x_\alpha\} = \{x_{++}, x_{+-}, x_{-+}, x_{--}\}$ corresponding to the two fundamental representations $L(\Lambda_i)$, $i = 1, 2$, of dimensions $D_1 = 5$ and $D_2 = 4$.

The realization for the Chevalley generators of $\mathfrak{so}_5$ on $\mathbf{S} \equiv \mathbb{C}[x_i, x_\alpha]$ can be chosen as

$$\begin{aligned}
 e_1 &= x_1 \frac{\partial}{\partial x_2} - x_{\bar{2}} \frac{\partial}{\partial x_{\bar{1}}} - x_{+-} \frac{\partial}{\partial x_{-+}}, \\
 e_2 &= \sqrt{2} \left(x_2 \frac{\partial}{\partial x_0} - x_0 \frac{\partial}{\partial x_{\bar{2}}} \right) - \left(x_{++} \frac{\partial}{\partial x_{+-}} - x_{-+} \frac{\partial}{\partial x_{--}} \right), \\
 h_1 &= x_1 \frac{\partial}{\partial x_1} - x_{\bar{1}} \frac{\partial}{\partial x_{\bar{1}}} - x_2 \frac{\partial}{\partial x_2} + x_{\bar{2}} \frac{\partial}{\partial x_{\bar{2}}} + x_{+-} \frac{\partial}{\partial x_{+-}} - x_{-+} \frac{\partial}{\partial x_{-+}}, \\
 h_2 &= 2 \left(x_2 \frac{\partial}{\partial x_2} - x_{\bar{2}} \frac{\partial}{\partial x_{\bar{2}}} \right) + \left(x_{++} \frac{\partial}{\partial x_{++}} + x_{-+} \frac{\partial}{\partial x_{-+}} - x_{+-} \frac{\partial}{\partial x_{+-}} - x_{--} \frac{\partial}{\partial x_{--}} \right), \\
 f_1 &= x_2 \frac{\partial}{\partial x_1} - x_{\bar{1}} \frac{\partial}{\partial x_{\bar{2}}} - x_{-+} \frac{\partial}{\partial x_{+-}}, \\
 f_2 &= \sqrt{2} \left(x_0 \frac{\partial}{\partial x_2} - x_{\bar{2}} \frac{\partial}{\partial x_0} \right) - \left(x_{+-} \frac{\partial}{\partial x_{++}} - x_{--} \frac{\partial}{\partial x_{-+}} \right).
 \end{aligned}
 \tag{2.36}$$

In this realization one finds the following explicit expressions for the metric g^{ij} (cf. (2.33))

$$g^{ij} = \frac{1}{2} \delta^{i\bar{j}}, \tag{2.37}$$

the γ -matrices $(\gamma^i)_\alpha{}^\beta$

$$\begin{aligned}
 \gamma^1 &= \tau^+ \otimes \tau^3, & \gamma^{\bar{1}} &= \tau^- \otimes \tau^3, \\
 \gamma^2 &= \tau^+ \otimes \tau^+, & \gamma^{\bar{2}} &= \tau^- \otimes \tau^-, \\
 \gamma^0 &= \sqrt{\frac{1}{2}} \tau^3 \otimes \tau^3,
 \end{aligned}
 \tag{2.38}$$

and the charge conjugation matrix $C^{\alpha\beta}$

$$C = \tau^1 \otimes i\tau^2, \tag{2.39}$$

where $\tau^\pm = \frac{1}{2}(\tau^1 \pm i\tau^2)$, and the τ^i , $i = 1, 2, 3$, are the standard Pauli matrices acting on the two-component coordinates $x_{\pm\pm}$ (see also section 4.4 for more explicit expressions).

Note that in the case of $\mathfrak{so}_5$, the third tensor product in (2.32) does not produce any additional relations, so we conclude that the ideal of quadratic relations, defining the

homogeneous coordinate ring $\mathbf{S}(V)$ of the (complete) flag variety for $\mathfrak{so}_5$, is generated by σ and σ_α of (2.33).

Now consider the following sequence of homomorphisms (see also Fig. 2.3)

$$(2.40) \quad 0 \rightarrow F^{(3)} \xrightarrow{d_3} F^{(2)} \xrightarrow{d_2} F^{(1)} \xrightarrow{d_1} F^{(0)} \rightarrow \mathbf{S}/I \rightarrow 0,$$

where

$$(2.41) \quad \begin{aligned} F^{(0)} &\cong \mathbf{S}(0, 0)_1, \\ F^{(1)} &\cong \mathbf{S}(-1, -1)_4 \oplus \mathbf{S}(-2, 0)_1, \\ F^{(2)} &\cong \mathbf{S}(-1, -2)_1 \oplus \mathbf{S}(-2, -1)_4, \\ F^{(3)} &\cong \mathbf{S}(-3, -2)_1, \end{aligned}$$

The homomorphisms d_i are defined by

$$(2.42) \quad \begin{aligned} d_1 : F^{(1)} &\rightarrow F^{(0)} : & e_\alpha^{(1)} &\mapsto \sigma_\alpha e^{(0)}, \\ & & e^{(1)} &\mapsto -\sigma e^{(0)}, \\ d_2 : F^{(2)} &\rightarrow F^{(1)} : & e^{(2)} &\mapsto x^\alpha e_\alpha^{(1)}, \\ & & e_\alpha^{(2)} &\mapsto x_\alpha e^{(1)} - (\gamma^i)_\alpha{}^\beta x_i e_\beta^{(1)}, \\ d_3 : F^{(3)} &\rightarrow F^{(2)} : & e^{(3)} &\mapsto \sigma e^{(2)} + \sigma^\alpha e_\alpha^{(2)}, \end{aligned}$$

where σ and σ_α are given in (2.33), and where $e_a^{(i)}$ denotes the generator of an $\mathbf{S}(a, b)_V$ component of $F^{(i)}$.

Using the properties (2.34) and (2.35) it is straightforward to verify that (2.40) defines a complex of $\mathfrak{so}_5$ modules. In fact, it is not too hard to show that it actually provides a resolution of $\mathbf{S}(V) = \mathbf{S}/I$.




Fig. 2.3. Resolution of $\mathbf{S}(V)$ for $\mathfrak{so}_5$

Applying the Euler-Poincaré principle to the resolution (2.40) we find an explicit formula for the character $h_V(M_1, M_2)$ of $L(M_1\Lambda_1 + M_2\Lambda_2)$, in particular, with (2.3),

$$(2.43) \quad \begin{aligned} \dim L(M_1\Lambda_1 + M_2\Lambda_2) = & \binom{M_1+4}{4} \binom{M_2+3}{3} - 4 \binom{M_1+3}{4} \binom{M_2+2}{3} \\ & - \binom{M_1+2}{4} \binom{M_2+3}{3} + \binom{M_1+3}{4} \binom{M_2+1}{3} \\ & + 4 \binom{M_1+2}{4} \binom{M_2+2}{3} - \binom{M_1+1}{4} \binom{M_2+1}{3}. \end{aligned}$$

3. AFFINIZED FLAG VARIETIES AND q -IDENTITIES

In [Bo] we introduced the concept of an affinized projective variety. Consider a projective variety V , defined by an ideal $I = \mathbf{I}(V) \subset \mathbb{C}[x_1, \dots, x_n]$, generated by a set of homogeneous elements f_i , $i = 1, \dots, t$. The affinized projective variety $\widehat{V}$ is the infinite dimensional projective variety defined by the ideal $\widehat{I} = \mathbf{I}(\widehat{V})$ generated by the relations $f_i[m]$, $i = 1, \dots, t$, $m \in \mathbb{Z}_{\geq 0}$, in $\widehat{\mathbf{S}} = \mathbb{C}[x_1[m], \dots, x_n[m]]_{m \in \mathbb{Z}_{\geq 0}}$, where the generators $f_i[m]$ of $\widehat{I}$ are obtained from f_i by replacing all monomials $x_{i_1} \dots x_{i_r}$ in f_i by

$$(3.1) \quad (x_{i_1} \dots x_{i_r})[m] = \sum_{\substack{n_{i_1}, \dots, n_{i_r} \geq 0 \\ n_{i_1} + \dots + n_{i_r} = m}} x_{i_1}[n_{i_1}] \dots x_{i_r}[n_{i_r}].$$

The homogeneous coordinate ring $\mathbf{S}(\widehat{V})$ of the affinized projective variety $\widehat{V}$ is graded both by the multi-degree inherited from the underlying projective variety V as well as the energy m , i.e.,

$$(3.2) \quad \deg(x_i[m]) = (\deg(x_i); m).$$

In the context of flag varieties, we have an additional grading corresponding to the $\mathfrak{g}$ -weight λ . Denoting by $\mathbf{S}(\widehat{V})_{(\mathbf{M}; N; \lambda)}$ the vector space of homogeneous polynomials f of multi-degree $(\mathbf{M}; N)$ and weight λ in $\mathbf{S}(\widehat{V})$, the (partial) Hilbert series of $\widehat{V}$ is defined as the (formal) $\mathfrak{g}$ -character

$$(3.3) \quad h_{\widehat{V}}(\mathbf{M}; q) = \sum_{N \geq 0, \lambda} \dim \mathbf{S}(\widehat{V})_{(\mathbf{M}; N; \lambda)} q^N e^\lambda.$$

The main result of [Bo] was the explicit computation of the (partial) Hilbert series $h_{\widehat{V}}(\mathbf{M}; q)$ for the coordinate ring of an affinized projective variety associated to a quadratic monomial ideal $I \subset \mathbf{S} \equiv \mathbb{C}[x_1, \dots, x_n]$. If $I = \langle x_i x_j \rangle_{(i,j) \in \mathcal{P}} \subset \mathbf{S}$ for some set $\mathcal{P}$ of (ordered)

pairs (i, j) , $i < j$, with $i, j \in \{1, \dots, n\}$, and $\mathbf{M} = (M_1, \dots, M_n)$ is the multi-degree where M_i denotes the number of x_i in a monomial, it was found that (see [Bo], section 4.1)

$$(3.4) \quad h_{\widehat{V}}(\mathbf{M}; q) = \frac{q^{\sum_{(i,j) \in \mathcal{P}} M_i M_j}}{(q)_{M_1} \dots (q)_{M_n}}.$$

In [Bo] we also gave an algorithm to compute an alternating sum expression for $h_{\widehat{V}}(\mathbf{M}; q)$ based on Taylor's resolution of $\mathbf{S}/I$. This algorithm is based on the identity

$$(3.5) \quad \frac{q^{M_1 M_2}}{(q)_{M_1} (q)_{M_2}} = \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m (q)_{M_1-m} (q)_{M_2-m}},$$

discovered and proved in [BS1, BS2] and shown to be related to the coordinate ring of the affinized projective variety associated to the ideal $I = \langle x_1 x_2 \rangle \subset \mathbb{C}[x_1, x_2]$ in [Bo].

Remark 3.1. The identity (3.5), as well as its ‘inverse’ (see [BS2])

$$(3.6) \quad \frac{1}{(q)_{M_1} (q)_{M_2}} = \sum_{m \geq 0} \frac{q^{(M_1-m)(M_2-m)}}{(q)_m (q)_{M_1-m} (q)_{M_2-m}},$$

are intimately related to, in fact can be used to prove, the five-term identity for Rogers’ dilogarithm [BCR].

In the remainder of this section we will explain how the results of [Bo], summarized above, can be used to compute the Hilbert series of affinized flag varieties.

As we have seen in section 2, the ideal that defines flag varieties, albeit quadratic, is not a monomial ideal. However, for the purpose of calculating the Hilbert series, we can replace I by the ideal of leading terms $\langle \text{LT}(I) \rangle$ with respect to any ordering on $\mathbf{S}$. Indeed, (cf. Ch. 9.3, Prop. 9 in [CLO1])

$$(3.7) \quad h_{\mathbf{S}/I}(\mathbf{M}; q) = h_{\mathbf{S}/\langle \text{LT}(I) \rangle}(\mathbf{M}; q).$$

We recall that $\langle \text{LT}(I) \rangle$ is the ideal generated by the leading terms $\text{LT}(f)$ for all $f \in I$. The ideal $\langle \text{LT}(I) \rangle$ is finitely generated (by Hilbert’s basis theorem) and a set of generators is given by $\text{LT}(g_i)$, $i = 1, \dots, s$, where $\{g_i\}_{i=1}^s$ is a Gröbnerbasis for I (see, e.g., [CLO1, CLO2]).

Unfortunately, by computing the Hilbert series by means of $\langle \text{LT}(I) \rangle$ we give up the manifest $\mathfrak{g}$ symmetry, since $\mathbf{S}/\langle \text{LT}(I) \rangle$ is no longer a $\mathfrak{g}$ -module. In examples we will see that the explicit $\mathfrak{g}$ -structure of the Hilbert series can be restored by successive application of (3.5).

Remark 3.2. In fact, note that the passing from I to $\langle \text{LT}(I) \rangle$ is reminiscent of taking the crystal limit of $U_q(\mathfrak{g})$ modules. It would be interesting to explore this connection further.

More seriously, while $\langle \text{LT}(I) \rangle$ is a monomial ideal, it is in general no longer quadratic (with the exception of $\mathfrak{sl}_3$, see section 4.2). However, non-quadratic monomial ideals may be transformed into quadratic monomial ideals by introducing additional variables. Having achieved this, we can then apply the results of [Bo] and derive an explicit expression for the Hilbert series $h_{\widehat{V}}(\mathbf{M}; q)$ of the affinized flag varieties $\widehat{V}$.

Let us illustrate in an example how the introduction of additional variables solves the problem above. Consider $I = \langle x_1 x_2 \dots x_n \rangle \subset \mathbb{C}[x_1, \dots, x_n]$. Clearly, for $n \geq 3$,

$$(3.8) \quad \mathbb{C}[x_1, \dots, x_n]/I \cong \mathbb{C}[x_1, \dots, x_n, t_1, \dots, t_{n-2}]/\widetilde{I},$$

where

$$(3.9) \quad \widetilde{I} = \langle t_1 - x_1 x_2, t_2 - t_1 x_3, \dots, t_{n-2} - t_{n-3} x_{n-1}, t_{n-2} x_n \rangle.$$

The isomorphism (3.8) is multi-degree preserving provided we assign

$$(3.10) \quad \deg(t_i) = \underbrace{(1, 1, \dots, 1)}_{i+1}, 0, \dots, 0).$$

With respect to the lexicographic ordering defined by

$$(3.11) \quad x_1 > x_2 > \dots > x_n > t_1 > \dots > t_{n-2},$$

we find

$$(3.12) \quad \langle \text{LT}(\widetilde{I}) \rangle = \langle x_1 x_2, t_1 x_3, \dots, t_{n-2} x_n \rangle,$$

which happens to be a quadratic monomial ideal. Thus, we can apply the results from [Bo] and conclude that for the corresponding affinized variety $\widehat{V}$ we have

$$(3.13) \quad h_{\widehat{V}}(\mathbf{M}; q) = \sum_{m_1, \dots, m_{n-2} \geq 0} \frac{q^Q}{\prod_{i=1}^n (q)_{M_i - \Delta M_i}},$$

where the quadratic form Q is given by

$$(3.14) \quad \begin{aligned} Q &= (M_1 - (m_1 + \dots + m_{n-2}))(M_2 - (m_1 + \dots + m_{n-2})) \\ &+ m_1(M_3 - (m_2 + \dots + m_{n-2})) + \dots + m_{n-2}M_n, \end{aligned}$$

and

$$(3.15) \quad \begin{aligned} \Delta M_1 &= m_1 + \dots + m_{n-2}, \\ \Delta M_2 &= m_1 + \dots + m_{n-2}, \\ \Delta M_3 &= m_2 + \dots + m_{n-2}, \\ &\vdots = \vdots \\ \Delta M_{n-1} &= m_{n-2}, \\ \Delta M_n &= 0. \end{aligned}$$

Note, moreover, that by repeated application of (3.5) and (3.6), eqn. (3.13) can be written as

$$(3.16) \quad h_{\widehat{V}}(M; q) = \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} \frac{1}{(q)_{M_1-m} \cdots (q)_{M_n-m}}.$$

This identity turns out to play an important role in the spinon description of $(\widehat{\mathfrak{sl}}_n)_{k=1}$ modules [BS1,BS2,BCR].

The above example illustrates that, for the purpose of computing the Hilbert series, non-quadratic monomial ideals can always be reduced to quadratic monomial ideals by the introduction of additional variables. For further examples we refer to section 4.

4. EXAMPLES

In this section we will explicitly compute the (partial) Hilbert series for the affinized flag varieties of $\mathfrak{sl}_n$, $n = 2, 3, 4$, and $\mathfrak{so}_{2n+1}$, $n = 2, 3$.

4.1. $\mathfrak{sl}_2$.

Recall from section 2.2.1 that the coordinate ring of the flag variety V for $\mathfrak{sl}_2$ is isomorphic to $\mathbf{S} = \mathbb{C}[x_1, x_2]$, i.e., the ideal of quadratic relations is trivial. Hence, the Hilbert series of the affinized variety $\widehat{V}$ is simply given by

$$(4.1) \quad h_{\widehat{V}}(M; q) = \sum_{m_1+m_2=M} \frac{1}{(q)_{m_1}(q)_{m_2}} e^{m_1\epsilon_1+m_2\epsilon_2},$$

which can obviously be interpreted as an ‘affinization’ of equation (2.19).

4.2. $\mathfrak{sl}_3$.

The coordinate ring of the flag variety V for $\mathfrak{sl}_3$ is given by $\mathbf{S}(V) = \mathbf{S}/I$ with $\mathbf{S} = \mathbb{C}[x_i, x_{\bar{i}}]_{i=1,2,3}$, and $I = \langle x_1x_{\bar{1}} + x_2x_{\bar{2}} + x_3x_{\bar{3}} \rangle$ (cf. section 2.2.2 where we used the notation $x^i = x_{\bar{i}}$). With respect to the lexicographic ordering defined by

$$(4.2) \quad x_1 > x_2 > x_3 > x_{\bar{1}} > x_{\bar{2}} > x_{\bar{3}},$$

we obviously have

$$(4.3) \quad \langle \text{LT}(I) \rangle = \langle x_1x_{\bar{1}} \rangle.$$

Thus, we can immediately apply our result for quadratic monomial ideals and conclude that the partial Hilbert series of the affinization $\widehat{V}$ is given by

$$(4.4) \quad h_{\widehat{V}}(M_1, M_2; q) = \sum_{\substack{m_1+m_2+m_3=M_1 \\ m_{\bar{1}}+m_{\bar{2}}+m_{\bar{3}}=M_2}} \frac{q^{m_1m_{\bar{1}}}}{\prod_i (q)_{m_i}(q)_{m_{\bar{i}}}} e^{\sum_i (m_i-m_{\bar{i}})\epsilon_i}.$$

As discussed in section 3, in order to use our results for quadratic monomial ideals, we have had to pass from I to $\langle \text{LT}(I) \rangle$, thereby breaking the manifest $\mathfrak{sl}_3$ symmetry. The result (4.4), however, can be written as an explicit $\mathfrak{sl}_3$ character by using (3.5), i.e.,

$$(4.5) \quad h_{\widehat{V}}(M_1, M_2; q) = \sum_{\substack{m_1+m_2+m_3+m=M_1 \\ m_{\bar{1}}+m_{\bar{2}}+m_{\bar{3}}+m=M_2}} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} \frac{1}{\prod_i (q)_{m_i} (q)_{m_{\bar{i}}}} e^{\sum_i (m_i - m_{\bar{i}}) \epsilon_i},$$

or, as

$$(4.6) \quad h_{\widehat{V}}(M_1, M_2; q) = \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} h_{\widehat{S}}(M_1 - m, M_2 - m; q),$$

where

$$(4.7) \quad h_{\widehat{S}}(M_1, M_2; q) = \sum_{\substack{m_1+m_2+m_3=M_1 \\ m_{\bar{1}}+m_{\bar{2}}+m_{\bar{3}}=M_2}} \frac{1}{\prod_i (q)_{m_i} (q)_{m_{\bar{i}}}} e^{\sum_i (m_i - m_{\bar{i}}) \epsilon_i},$$

denotes the Hilbert series of the affinization of the coordinate ring $\mathbf{S} = \mathbb{C}[x_i, x_{\bar{i}}]$. Clearly, (4.7) has to be interpreted as the Hilbert series obtained by applying the Euler-Poincaré principle to an affinization of the resolution (2.22). In this case the affinization of the resolution is easily constructed and yields a Koszul complex (cf. [Bo]).

4.3. $\mathfrak{sl}_4$.

The coordinate ring for the flag variety of $\mathfrak{sl}_4$ is given by $I \subset \mathbb{C}[x_i, x_{ij}, x_{ijk}] = \mathbf{S}$ where the generators of I are explicitly given in section 2.2.3. With respect to the lexicographic ordering on $\mathbf{S}$ defined by

$$(4.8) \quad \begin{aligned} x_1 &> x_2 > x_3 > x_4 > \\ x_{12} &> x_{13} > x_{14} > x_{23} > x_{24} > x_{34} > \\ x_{123} &> x_{124} > x_{134} > x_{234}, \end{aligned}$$

we find

$$(4.9) \quad \begin{aligned} \langle \text{LT}(I) \rangle &= \langle x_1 x_{23}, x_1 x_{24}, x_1 x_{34}, x_1 x_{234}, x_2 x_{34}, x_2 x_{13} x_{24}, x_{12} x_{34}, \\ &\quad x_{12} x_{134}, x_{12} x_{234}, x_{13} x_{234}, x_{13} x_{24} x_{134}, x_{14} x_{234} \rangle. \end{aligned}$$

Note that in this case the ideal of leading terms $\langle \text{LT}(I) \rangle$ is not a quadratic monomial ideal. We can however apply the trick discussed in section 3 and introduce an additional variable

$$(4.10) \quad t = x_{13} x_{24},$$

of multi-degree $\deg(t) = (0, 2, 0)$ and vanishing $\mathfrak{sl}_4$ -weight. Indeed,

$$(4.11) \quad \mathbb{C}[x_i, x_{ij}, x_{ijk}]/\langle \text{LT}(I) \rangle \cong \mathbb{C}[x_i, x_{ij}, x_{ijk}, t]/\tilde{I},$$

where

$$(4.12) \quad \begin{aligned} \tilde{I} = \langle & x_1x_{23}, x_1x_{24}, x_1x_{34}, x_1x_{234}, x_2x_{34}, x_2t, x_{12}x_{34}, \\ & x_{12}x_{134}, x_{12}x_{234}, x_{13}x_{234}, tx_{134}, x_{14}x_{234}, t - x_{13}x_{24} \rangle. \end{aligned}$$

Extending the lexicographic ordering on $\mathbf{S}$ defined by (4.8) to $\tilde{\mathbf{S}} = \mathbb{C}[x_i, x_{ij}, x_{ijk}, t]$ by

$$(4.13) \quad x_1 > \dots > x_{234} > t,$$

we can now pass to the ideal of leading terms of $\tilde{I}$. We find

$$(4.14) \quad \begin{aligned} \langle \text{LT}(\tilde{I}) \rangle = \langle & x_1x_{23}, x_1x_{24}, x_1x_{34}, x_1x_{234}, x_1t, x_2x_{34}, x_2t, x_{12}x_{34}, \\ & x_{12}x_{134}, x_{12}x_{234}, x_{13}x_{24}, x_{13}x_{234}, x_{14}x_{234}, x_{134}t, x_{234}t \rangle. \end{aligned}$$

Thus, $\langle \text{LT}(\tilde{I}) \rangle$ is a quadratic monomial ideal, and we are finally in a position to apply the results of [Bo]. We find

$$(4.15) \quad \begin{aligned} h_{\hat{\mathbf{V}}}(M_1, M_2, M_3; q) = & \sum_{\substack{\sum m_i = M_1 \\ \sum m_{ij} + 2m = M_2 \\ \sum m_{ijk} = M_3}} \frac{q^Q}{(q)_m \prod_i (q)_{m_i} \prod_{i < j} (q)_{m_{ij}} \prod_{i < j < k} (q)_{m_{ijk}}} \\ & \times e^{\sum m_i \epsilon_i + \sum m_{ij}(\epsilon_i + \epsilon_j) + \sum m_{ijk}(\epsilon_i + \epsilon_j + \epsilon_k)}, \end{aligned}$$

with quadratic form

$$(4.16) \quad \begin{aligned} Q = & m_1(m_{23} + m_{24} + m_{34} + m_{234} + m) + m_2(m_{34} + m) + m_{12}m_{34} \\ & + m_{13}m_{24} + m_{134}(m_{12} + m) + m_{234}(m_{12} + m_{13} + m_{14} + m). \end{aligned}$$

As in section 4.2 we may now attempt to rewrite (4.15) as a manifest $\mathfrak{sl}_4$ character by successively applying (3.5) according to the algorithm outlined in [Bo]. One would expect that the result can be written in a form that can be interpreted as arising from a properly affinized version of the resolution of Fig. 2.2. We have not succeeded in carrying out this program in all generality, but partial results (relevant to the applications discussed in [BS4]) are easily obtained, e.g., (cf. (4.6))

$$(4.17) \quad h_{\hat{\mathbf{V}}}(M_1, 0, M_3; q) = \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} h_{\hat{\mathbf{S}}}(M_1 - m, 0, M_3 - m; q),$$

where

$$(4.18) \quad h_{\hat{\mathbf{S}}}(M_1, 0, M_3; q) = \sum_{\substack{\sum m_i = M_1 \\ \sum m_{ijk} = M_3}} \frac{1}{\prod_i (q)_{m_i} \prod_{i < j < k} (q)_{m_{ijk}}} e^{\sum_i m_i \epsilon_i + \sum m_{ijk} \epsilon_{ijk}},$$

denotes the Hilbert series of the affinization of the coordinate ring $\mathbf{S} = \mathbb{C}[x_i, x_{ijk}]$. The result (4.18) generalizes in an obvious way to $\mathfrak{sl}_n$.

4.4. $\mathfrak{so}_5$.

For $\mathfrak{so}_5$ we need to consider $I \subset \mathbb{C}[x_i, x_\alpha] = \mathbf{S}$ where $I = \langle \sigma, \sigma_\alpha \rangle$ is given in section 2.3.1. Explicitly,

$$\begin{aligned}
 \sigma &= x_1 x_{\bar{1}} + x_2 x_{\bar{2}} + \frac{1}{2} x_0 x_0, \\
 \sigma_{++} &= x_1 x_{-+} + \sqrt{\frac{1}{2}} x_0 x_{++} + x_2 x_{+-}, \\
 \sigma_{+-} &= -x_1 x_{--} - \sqrt{\frac{1}{2}} x_0 x_{+-} + x_2 x_{++}, \\
 \sigma_{-+} &= x_2 x_{--} - \sqrt{\frac{1}{2}} x_0 x_{-+} + x_{\bar{1}} x_{++}, \\
 \sigma_{--} &= x_{\bar{2}} x_{-+} + \sqrt{\frac{1}{2}} x_0 x_{--} - x_{\bar{1}} x_{+-}.
 \end{aligned}
 \tag{4.19}$$

The (minimal) Gröbnerbasis with respect to the lexicographic ordering on $\mathbf{S}$ defined by

$$x_1 > x_2 > x_0 > x_{\bar{2}} > x_{\bar{1}} > x_{++} > x_{+-} > x_{-+} > x_{--},
 \tag{4.20}$$

is given by $\{\sigma, \sigma_\alpha, \tau\}$ where

$$\begin{aligned}
 \tau &= -\sqrt{\frac{1}{2}} x_0 x_{\bar{1}} x_{++} - x_2 x_{\bar{1}} x_{+-} + \frac{1}{2} x_0 x_0 x_{-+} + x_2 x_{\bar{2}} x_{-+} \\
 &= \sigma x_{-+} - x_{\bar{1}} \sigma_{++}.
 \end{aligned}
 \tag{4.21}$$

The monomial ideal $\langle \text{LT}(I) \rangle$ generated by the leading terms in I is therefore given by

$$\langle \text{LT}(I) \rangle = \langle x_1 x_{\bar{1}}, x_1 x_{-+}, x_1 x_{--}, x_2 x_{\bar{2}} x_{-+}, x_2 x_{--}, x_0 x_{--} \rangle.
 \tag{4.22}$$

Again, to apply the results of [Bo], we need to introduce one additional variable $t = x_2 x_{\bar{2}}$, of multi-degree $\deg(t) = (2, 0)$ and vanishing $\mathfrak{so}_5$ weight. Then

$$\mathbb{C}[x_i, x_\alpha] / \langle \text{LT}(I) \rangle \cong \mathbb{C}[x_i, x_\alpha, t] / \tilde{I},
 \tag{4.23}$$

with

$$\tilde{I} = \langle x_1 x_{\bar{1}}, x_1 x_{-+}, x_1 x_{--}, x_2 x_{--}, x_0 x_{--}, t - x_2 x_{\bar{2}}, t x_{-+} \rangle.
 \tag{4.24}$$

Extending the lexicographic ordering on $\mathbf{S}$ defined by (4.20) to $\tilde{\mathbf{S}} = \mathbb{C}[x_i, x_\alpha, t]$ by

$$x_1 > x_2 > x_0 > x_{\bar{2}} > x_{\bar{1}} > x_{++} > x_{+-} > x_{-+} > x_{--} > t,
 \tag{4.25}$$

we obtain

$$\langle \text{LT}(\tilde{I}) \rangle = \langle x_1 x_{\bar{1}}, x_1 x_{-+}, x_1 x_{--}, x_2 x_{\bar{2}}, x_2 x_{--}, x_0 x_{--}, x_{-+} t, x_{--} t \rangle,
 \tag{4.26}$$

which is a quadratic monomial ideal. Thus, we conclude that the Hilbert series of the coordinate ring of the affinized flag variety of $\mathfrak{so}_5$ is given by

$$(4.27) \quad h_{\widehat{V}}(M_1, M_2; q) = \sum_{\substack{\sum m_i + 2m = M_1 \\ \sum m_\alpha = M_2}} \frac{q^Q}{\prod_i (q)_{m_i} \prod_\alpha (q)_{m_\alpha}} e^{\sum m_i \lambda_i + \sum m_\alpha \lambda_\alpha},$$

where

$$(4.28) \quad Q = m_1(m_{\bar{1}} + m_{-+} + m_{--}) + m_2(m_{\bar{2}} + m_{--}) + m_0 m_{--} + m(m_{-+} + m_{--}),$$

and $\{\lambda_i\}$ and $\{\lambda_\alpha\}$ denote the weights of the representations $L(\Lambda_1)$ and $L(\Lambda_2)$, respectively (see section 2.3).

Again, in principle one could proceed to write (4.27) as a manifest $\mathfrak{so}_5$ character by successive application of (3.5). We have not been able to carry this out in all generality, but the following partial results can be proved

$$(4.29) \quad \begin{aligned} h_{\widehat{V}}(0, M_2; q) &= h_{\widehat{S}}(0, M_2; q), \\ h_{\widehat{V}}(M_1, 0; q) &= \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} h_{\widehat{S}}(M_1 - 2m, 0; q), \\ h_{\widehat{V}}(M_1, 1; q) &= \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} h_{\widehat{S}}(M_1 - 2m, 1; q) \\ &\quad - \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m+1)}}{(q)_m (q)_1} h_{\widehat{S}}(M_1 - 2m - 1, 0; q) \chi_4 \\ &\quad + \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m+1)}}{(q)_m (q)_1} h_{\widehat{S}}(M_1 - 2m - 2, 0; q) \chi_4, \end{aligned}$$

where

$$(4.30) \quad h_{\widehat{S}}(M_1, M_2; q) = \sum_{\substack{\sum m_i = M_1 \\ \sum m_\alpha = M_2}} \frac{1}{\prod_i (q)_{m_i} \prod_\alpha (q)_{m_\alpha}} e^{\sum m_i \lambda_i + \sum m_\alpha \lambda_\alpha},$$

denotes the Hilbert series of the affinization of $\mathbf{S} = \mathbb{C}[x_i, x_\alpha]$. The third equation can be simplified, but we have left it in this form to elucidate its origin as arising from an ‘affinization’ of the resolution (2.40) (see also Fig. 2.3).

4.5. $\mathfrak{so}_7$.

For $\mathfrak{so}_7$ we only quote the result for the coordinate ring of the partial flag variety discussed in section 2.3. The procedure is as before, so we will only give the main intermediate

steps that lead to the Hilbert series. We use the notation of section 2.3. With respect to the lexicographic ordering defined by

$$(4.31) \quad \begin{aligned} & x_1 > x_2 > x_3 > x_0 > x_{\bar{3}} > x_{\bar{2}} > x_{\bar{1}} > \\ & x_{+++} > x_{++-}x_{-+-} > x_{-++} > x_{+--} > x_{-+-} > x_{--+} > x_{---}, \end{aligned}$$

we have

$$(4.32) \quad \begin{aligned} \langle \text{LT}(I) \rangle = & \langle x_1x_{\bar{1}}, x_{+++}x_{---}, x_1x_{-++}, x_1x_{-+-}, x_1x_{--+}, x_1x_{---}, x_2x_{--+}, \\ & x_2x_{---}, x_3x_{---}, x_0x_{---}, x_2x_{\bar{2}}x_{-++}, x_2x_{\bar{2}}x_{-+-}, x_2x_{+-+}x_{-+-}, \\ & x_3x_{\bar{3}}x_{--+}, x_3x_{++-}x_{--+}, x_0x_{++-}x_{--+}, x_3x_{\bar{3}}x_{+-+}x_{-+-} \rangle. \end{aligned}$$

To transform this back into a quadratic monomial ideal we need to introduce four additional variables for which one can take

$$(4.33) \quad \begin{aligned} t_1 &= x_{++-}x_{--+}, & t_2 &= x_{+-+}x_{-+-}, \\ t_3 &= x_3x_{\bar{3}}, & t_4 &= x_2x_{\bar{2}}. \end{aligned}$$

Obviously, the variables (4.33) all have vanishing $\mathfrak{so}_7$ -weight and multi-degrees

$$(4.34) \quad \begin{aligned} \deg(t_1) &= \deg(t_2) = (0, 0, 2), \\ \deg(t_3) &= \deg(t_4) = (2, 0, 0). \end{aligned}$$

We have

$$(4.35) \quad \mathbb{C}[x_i, x_\alpha] / \langle \text{LT}(I) \rangle \cong \mathbb{C}[x_i, x_\alpha, t_1, t_2, t_3, t_4] / \tilde{I},$$

where

$$(4.36) \quad \begin{aligned} \tilde{I} = & \langle x_1x_{\bar{1}}, x_{+++}x_{---}, x_1x_{-++}, x_1x_{-+-}, x_1x_{--+}, x_1x_{---}, x_2x_{--+}, x_2x_{---}, \\ & x_3x_{---}, x_0x_{---}, t_1 - x_{++-}x_{--+}, t_2 - x_{+-+}x_{-+-}, t_3 - x_3x_{\bar{3}}, \\ & t_4 - x_2x_{\bar{2}}, t_1x_3, t_1x_0, t_2x_2, t_3x_{--+}, t_2t_3, t_4x_{-+-}, t_4x_{--+} \rangle. \end{aligned}$$

Extending the ordering defined by (4.31) by

$$(4.37) \quad x_1 > \dots > x_{---} > t_1 > t_2 > t_3 > t_4,$$

we find

$$(4.38) \quad \begin{aligned} \langle \text{LT}(\tilde{I}) \rangle = & \langle x_1x_{\bar{1}}, x_1x_{-++}, x_1x_{-+-}, x_1x_{--+}, x_1x_{---}, x_1t_1, x_1t_2, \\ & x_2x_{\bar{2}}, x_2x_{--+}, x_2x_{---}, x_2t_1, x_2t_2, \\ & x_3x_{\bar{3}}, x_3x_{---}, x_3t_1, x_0x_{---}, x_0t_1, \\ & x_{+++}x_{---}, x_{++-}x_{--+}, x_{+-+}x_{-+-}, \\ & x_{-++}t_4, x_{-+-}t_4, x_{--+}t_3, x_{--+}t_4, x_{---}t_3, x_{---}t_4, \\ & t_1t_3, t_1t_4, t_2t_3, t_2t_4 \rangle. \end{aligned}$$

Thus, the result for the Hilbert series is

$$(4.39) \quad h_{\widehat{V}}(M_1, 0, M_3; q) = \sum_{\substack{\sum m_i + 2p_3 + 2p_4 = M_1 \\ \sum m_\alpha + 2p_1 + 2p_2 = M_3}} \frac{q^Q}{\prod_i (q)_{m_i} \prod_\alpha (q)_{m_\alpha} \prod_j (q)_{p_j}} \times e^{\sum m_i \lambda_i + \sum m_\alpha \lambda_\alpha},$$

with

$$(4.40) \quad \begin{aligned} Q = & m_1(m_{\bar{1}} + m_{-++} + m_{-+-} + m_{--+} + m_{---} + p_1 + p_2) \\ & + m_2(m_{\bar{2}} + m_{--+} + m_{---} + p_1 + p_2) \\ & + m_3(m_{\bar{3}} + m_{---} + p_1) + m_0(m_{---} + p_1) \\ & + m_{+++} + m_{---} + m_{++-} + m_{--+} + m_{+-+} + m_{-+-} \\ & + (m_{-++} + m_{-+-})p_4 + (m_{--+} + m_{---})(p_3 + p_4) \\ & + (p_1 + p_2)(p_3 + p_4). \end{aligned}$$

As in (4.29) we can write the character (4.39) as a manifest $\mathfrak{so}_7$ character in the following cases

$$(4.41) \quad \begin{aligned} h_{\widehat{V}}(0, 0, M_3; q) &= \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} h_{\widehat{\mathbf{S}}}(0, 0, M_3 - 2m; q), \\ h_{\widehat{V}}(M_1, 0, 0; q) &= \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} h_{\widehat{\mathbf{S}}}(M_1 - 2m, 0, 0; q), \\ h_{\widehat{V}}(M_1, 0, 1; q) &= \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m-1)}}{(q)_m} h_{\widehat{\mathbf{S}}}(M_1 - 2m, 0, 1; q) \\ &\quad - \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m+1)}}{(q)_m (q)_1} h_{\widehat{\mathbf{S}}}(M_1 - 2m - 1, 0, 0; q) \chi_{\mathbf{8}} \\ &\quad + \sum_{m \geq 0} (-1)^m \frac{q^{\frac{1}{2}m(m+1)}}{(q)_m (q)_1} h_{\widehat{\mathbf{S}}}(M_1 - 2m - 2, 0, 0; q) \chi_{\mathbf{8}}, \end{aligned}$$

where

$$(4.42) \quad h_{\widehat{\mathbf{S}}}(M_1, 0, M_3; q) = \sum_{\substack{\sum m_i = M_1 \\ \sum m_\alpha = M_3}} \frac{1}{\prod_i (q)_{m_i} \prod_\alpha (q)_{m_\alpha}} e^{\sum m_i \lambda_i + \sum m_\alpha \lambda_\alpha},$$

denotes the Hilbert series of the affinization of $\mathbf{S} = \mathbb{C}[x_i, x_\alpha]$. The second and third of these identities have an obvious generalization to $\mathfrak{so}_{2n+1}$, while the generalization of the first will be more involved due to additional relations arising from the tensor product $\text{Sym}^2 L(\Lambda_n)$ (see (2.32)).

5. RELATION TO MODIFIED HALL-LITTLEWOOD
POLYNOMIALS AND AFFINE LIE ALGEBRA CHARACTERS

By construction, $P(\mathbf{M}; q) \equiv \left(\prod_{i=1}^{\ell} (q)_{M_i} \right) h_{\widehat{V}}(\mathbf{M}; q)$ is a $\mathfrak{g}$ -character valued polynomial in q such that

$$(5.1) \quad \lim_{q \rightarrow 1} P(\mathbf{M}; q) = \chi_{\Lambda_1}^{M_1} \cdots \chi_{\Lambda_{\ell}}^{M_{\ell}}.$$

In other words, $P(\mathbf{M}; q)$ is a q -deformation of the character of the tensor product $L(\Lambda_1)^{\otimes M_1} \otimes \cdots \otimes L(\Lambda_{\ell})^{\otimes M_{\ell}}$. A natural q -deformation, that shows up in many contexts, is the so-called modified Hall-Littlewood polynomial (see [Ki1, Ki2] for reviews and recent results). For a general finite dimensional simple Lie algebra $\mathfrak{g}$ of rank ℓ , it can be defined as follows [KR]. Let Λ_i , α_i and $\alpha_i^{\vee}$ ($i = 1, \dots, \ell$) be the fundamental weights, simple roots and simple co-roots, respectively. For any pair of dominant integral weights λ, μ , define the polynomial $M_{\lambda\mu}(q)$ by

$$(5.2) \quad M_{\lambda\mu}(q) = \sum_{\mathbf{m}} q^{\widetilde{c}(\mathbf{m})} \prod_{i=1}^{\ell} \prod_{a=1}^{\infty} \left[\begin{matrix} P_a^{(i)}(\mathbf{m}) + m_a^{(i)} \\ m_a^{(i)} \end{matrix} \right],$$

where the sum is taken over all nonnegative integers $m_a^{(i)}$ ($i = 1, \dots, \ell$, $a = 1, 2, \dots$), such that

$$(5.3) \quad \mu - \lambda = \sum_{i=1}^{\ell} \left(\sum_{a=1}^{\infty} a m_a^{(i)} \right) \alpha_i.$$

Moreover,

$$(5.4) \quad P_a^{(i)}(\mathbf{m}) = (\mu, \alpha_i^{\vee}) - \sum_{j=1}^{\ell} \sum_{b=1}^{\infty} \Phi_{ab}^{ij} m_b^{(j)},$$

where

$$(5.5) \quad \Phi_{ab}^{ij} = \frac{2(\alpha_i, \alpha_j)}{\alpha_i^2 \alpha_j^2} \min(a\alpha_i^2, b\alpha_j^2),$$

and $\widetilde{c}(\mathbf{m})$ is the cocharge

$$(5.6) \quad \widetilde{c}(\mathbf{m}) = \frac{1}{2} \sum_{i,j=1}^{\ell} \sum_{a,b=1}^{\infty} m_a^{(i)} \Phi_{ab}^{ij} m_b^{(j)}.$$

For $\mathfrak{sl}_n$, the polynomials (5.2) are related to the Kostka-Foulkes polynomials $K_{\lambda\mu}(q)$ in a simple way (see, e.g., [Ki1]).

For λ dominant integral, let χ_λ denote the character of the finite dimensional irreducible $\mathfrak{g}$ -module $L(\lambda)$. Then, for dominant integral weight λ , the modified Hall-Littlewood polynomial $M_\lambda(q)$ is defined as the character valued polynomial

$$(5.7) \quad M_\lambda(q) = \sum_{\mu} M_{\mu\lambda}(q) \chi_{\mu}.$$

It has the property that for $q = 1$ it equals the character of the tensor product module $W_1^{M_1} \otimes \dots \otimes W_\ell^{M_\ell}$, where $M_i = (\lambda, \alpha_i^\vee)$, and where W_i denotes the “minimal affinization” of $L(\Lambda_i)$, i.e., the minimal irreducible module of the quantum affine algebra $U_q(\widehat{\mathfrak{g}})$ (or Yangian $Y(\mathfrak{g})$) such that $L(\Lambda_i) \subset W_i$ [KR]. It is therefore natural to make the following conjecture

Conjecture 5.1. *Let $\lambda = M_1\Lambda_1 + \dots + M_\ell\Lambda_\ell$, where $M_i \geq 0$ for those i such that $W_i \cong L(\Lambda_i)$ (as a $\mathfrak{g}$ -module) and $M_i = 0$ for the remaining i . Then*

$$(5.8) \quad M_\lambda(q) = \left(\prod_{i=1}^{\ell} (q)_{M_i} \right) h_{\widehat{V}}(M_1, \dots, M_\ell; q).$$

Note that the condition $W_i \cong L(\Lambda_i)$ is satisfied for all i in the case of $\mathfrak{sl}_n$, and for $i = 1, n$, for $\mathfrak{so}_{2n+1}$, i.e., for all the examples discussed in this paper. We have checked Conjecture 5.1 numerically to high order for the examples in section 4 and have a proof in special cases for which other concise formulas for $M_\lambda(q)$ are known (see, in particular, [Ki1]). Special cases of this conjecture have also appeared in [BS4]. A general proof would clearly require a better understanding of the geometry of affinized flag varieties. It is conceivable that the conjecture can be lifted to all possible choices of M_i by repeating the analysis of this paper to the affinized coordinate ring of a variety defined by an ideal in $\mathbb{C}[x_a^{(i)}]$ where the coordinates now correspond to a basis of the $U_q(\widehat{\mathfrak{g}})$ -module W_i , rather than $L(\Lambda_i)$.

A different, but possibly related, relation between modified Hall-Littlewood polynomials and the geometry of flag varieties was observed in [HS,Sh] (see also [Ki2]).

The results of this paper can also be used to obtain explicit quasi-particle type expressions for the characters of the integrable highest weight modules of affine Lie algebras $\widehat{\mathfrak{g}}$, in particular of $\widehat{\mathfrak{sl}_n}$ and $\widehat{\mathfrak{so}_5}$ in view of our results in section 4. These characters can be written as [NY2,Ya,HKKOTY]

$$(5.9) \quad \text{ch}_\lambda(q) = \sum_{\mu=M_1\Lambda_1+\dots+M_\ell\Lambda_\ell} \frac{1}{\prod_{i=1}^{\ell} (q)_{M_i}} M_{\lambda\mu}^{(k)}(q) M_\mu(q),$$

where λ is the highest weight of the integrable $\widehat{\mathfrak{g}}$ -module and $M_{\lambda\mu}^{(k)}(q)$ a certain ‘level- k restriction’ of the polynomials (5.2) (see [BMS,DKKMM,Ki1,HKKOTY], and references therein, for explicit expressions in the case of $\mathfrak{sl}_n$, and [BS4] for $\mathfrak{so}_5$). In fact, by using (5.9), the characters obtained this way will be of the “Universal Chiral Partition Function” (UCPF) form which was recently argued to be a universal expression for the (chiral) characters of any conformal field theory [BM]. This form reads

$$(5.10) \quad \text{ch}_\lambda(q) = \sum_{\substack{m_1, \dots, m_n \geq 0 \\ \text{restrictions}}} q^{\frac{1}{2}\mathbf{m} \cdot \mathbf{G} \cdot \mathbf{m} - \frac{1}{2}\mathbf{A} \cdot \mathbf{m}} \prod_a \left[\frac{((\mathbf{1} - \mathbf{G}) \cdot \mathbf{m} + \frac{\mathbf{u}}{2})_a}{m_a} \right],$$

where $\mathbf{G}$ is an $n \times n$ matrix and $\mathbf{A}$ and $\mathbf{u}$ are certain n -vectors. Both $\mathbf{A}$ and $\mathbf{u}$ as well as the restrictions on the summations over the quasi-particle numbers m_a will in general depend on the sector λ , while $\mathbf{G}$ will be independent of λ .

Another interesting connection of modified Hall-Littlewood polynomials to affine Lie algebra characters was observed (and proved in special cases), in [Ki1,NY1,KMOTU,KKN,HKKOTY,BS4] for $\mathfrak{sl}_n$ and in [BS4] for $\mathfrak{so}_n$. It turns out that these characters can often be obtained from $M_\lambda(q^{-1})$ in a ‘large λ -limit’, i.e., in a limit where one of the Dynkin indices of λ tends to infinity while the others are kept fixed. This observation is intimately related to taking the TBA limit in the integrable spin chain underlying the definition of $M_\lambda(q)$ [KR].

The connection of affinized projective varieties with quasi-particles in conformal field theory, which in fact motivated this research, arises as follows. Quasi-particles in conformal field theory correspond to intertwiners (‘Chiral Vertex Operators’ or CVOs) between modules of the Chiral Algebra. The degrees of freedom in these CVOs can be separated, at least heuristically, in terms of pseudo-particles that generate the collection of possible ‘fusion paths’ for the CVOs (i.e., the sequence of modules between which the CVOs intertwine) and physical particles whose degrees of freedom can be interpreted as the coordinate ring of an affinized projective variety; the ideal corresponds precisely to the null-states in the physical quasi-particle Fock space. [In (5.10) pseudo-particles correspond to $u_a < \infty$ while physical particles have $u_a = \infty$.] This paper thus gives an effective technique to compute the contribution of the physical particles to the conformal field theory characters. It is hoped that similar techniques may be applied to extract the pseudo-particle contribution. We refer to [BCR] for a more detailed exposition of this connection, a more explicit discussion of the UCPF form of the various affine Lie algebra characters and an application to the exclusion statistics satisfied by these quasi-particles. The results of [BCR] can be taken as further evidence for the correctness of Conjecture 5.1.

ACKNOWLEDGEMENTS

P.B. is supported by a QEII research fellowship from the Australian Research Council. The various Gröbnerbasis computations were performed with the help of MathematicaTM.

REFERENCES

- [BM] A. Berkovich and B. McCoy, *The universal chiral partition function for exclusion statistics*, [hep-th/9808013].
- [BMS] A. Berkovich, B. McCoy and A. Schilling, *Rogers-Schur-Ramanujan type identities for the $M(p, p')$ minimal models of conformal field theory*, Comm. Math. Phys. **191** (1998), 325–395, [q-alg/9607020].
- [Bo] P. Bouwknegt, *q -Identities and affinized projective varieties, I. Quadratic monomial ideals*, [math-ph/9902010].
- [BCR] P. Bouwknegt, L.-H. Chim and D. Ridout, *Exclusion statistics in conformal field theory and the UCPF for WZW models*, preprint ADP-99-12/M78, in preparation.
- [BS1] P. Bouwknegt and K. Schoutens, *The $\widehat{SU(n)}_1$ WZW models: Spinon decomposition and Yangian structure*, Nucl. Phys. **B482** (1996), 345–372, [hep-th/9607064].
- [BS2] ———, *Spinon decomposition and Yangian structure of $\widehat{\mathfrak{sl}}_n$ modules*, Geometric Analysis and Lie Theory in Mathematics and Physics, Australian Mathematical Society Lecture Series **11** (A.L. Carey and M.K. Murray, eds.), Cambridge University Press, Cambridge, 1997, pp. 105–131, [q-alg/9703021].
- [BS3] ———, *Non-abelian electrons: $SO(5)$ superspin regimes for correlated electrons on a two-leg ladder*, Phys. Rev. Lett., to appear, [cond-mat/9805232].
- [BS4] ———, *Exclusion statistics in conformal field theory – generalized fermions and spinons for level-1 WZW models*, Nucl. Phys. **B**, to appear, [hep-th/9810113].
- [CLO1] D. Cox, J. Little and D. O’Shea, *Ideals, varieties, and algorithms: An introduction to computational algebraic geometry and commutative algebra*, Springer Verlag, Berlin, 1997.
- [CLO2] ———, *Using algebraic geometry*, Springer Verlag, Berlin, 1998.
- [DKKMM] S. Dasmahapatra, R. Kedem, T. Klassen, B. McCoy and E. Melzer, *Quasi-Particles, Conformal Field Theory, and q -Series*, Int. J. Mod. Phys. **B7** (1993), 3617–3648, [hep-th/9303013].
- [FS1] B.L. Feigin and A.V. Stoyanovsky, *Quasi-particles models for the representations of Lie algebras and geometry of flag manifold*, [hep-th/9308079].
- [FS2] ———, *Functional models for representations of current algebras and semi-infinite Schubert cells*, Funct. Anal. and Appl. **28** (1994), 55–72.
- [Fu] W. Fulton, *Young tableaux*, London Math. Soc. Student Texts **35**, Cambridge University Press, Cambridge, 1997.
- [FH] W. Fulton and J. Harris, *Representation theory; A first course*, Graduate Texts in Math. 129, Springer Verlag, Berlin, 1991.
- [GS] S. Guruswamy and K. Schoutens, *Non-abelian exclusion statistics*, [cond-mat/9903045].
- [HKKOTY] G. Hatayama, A. Kirillov, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, *Character formulae of $\widehat{\mathfrak{sl}}_n$ -modules and inhomogeneous paths*, Nucl. Phys. **B536** (1999), 575–616, [math.QA/9802085].
- [Ha] D. Haldane, “*Fractional statistics*” in arbitrary dimensions: a generalization of the Pauli principle, Phys. Rev. Lett. **67** (1991), 937–940.
- [HS] R. Hotta and N. Shimomura, *The fixed point subvarieties of unipotent transformations on generalized flag varieties and Green functions*, Math. Ann. **241** (1979), 193–208.
- [Ki1] A. Kirillov, *Dilogarithm identities*, Prog. Theor. Phys. Suppl. **118** (1995), 61–142, [hep-th/9408113].
- [Ki2] ———, *New combinatorial formula for modified Hall-Littlewood polynomials*, [math.QA/9803006].
- [KKN] A. Kirillov, A. Kuniba and T. Nakanishi, *Skew Young diagram method in spectral decomposition of integrable lattice models II: Higher levels*, Nucl. Phys. **B529** (1998), 611–638, [q-alg/9711009].
- [KR] A. Kirillov and N. Reshetikhin, *The Bethe ansatz and the combinatorics of Young tableaux*, J. Sov. Math. **41** (1988), 925–955.

- [KMOTU] A. Kuniba, K. Misra, M. Okado, T. Takagi and J. Uchiyama, *Demazure crystals and symmetric functions*, [q-**alg**/9612018].
- [LT] G. Lancaster and J. Towber, *Representation-functors and flag-algebras for the classical groups I*, J. Algebra **59** (1979), 16–38.
- [NY1] A. Nakayashiki and Y. Yamada, *Kostka polynomials and energy functions in solvable lattice models*, Selecta Math. (N.S.) **3** (1997), 547–599, [q-**alg**/9512027].
- [NY2] ———, *On spinon character formulas*, Frontiers in Quantum Field Theories (H. Itoyama et al., eds.), World Scientific, Singapore, 1996, pp. 367–371.
- [Sh] N. Shimomura, *A theorem of the fixed point set of a unipotent transformation of the flag manifold*, J. Math. Soc. Japan **32** (1980), 55–64.
- [Ya] Y. Yamada, *On q -Clebsch Gordan rules and the spinon character formulas for affine $C_2^{(1)}$ algebra*, [q-**alg**/9702019].

DEPARTMENT OF PHYSICS AND MATHEMATICAL PHYSICS, UNIVERSITY OF ADELAIDE, ADELAIDE
SA 5005, AUSTRALIA

E-mail address: pbouwkn@physics.adelaide.edu.au

DEPARTMENT OF PHYSICS AND MATHEMATICAL PHYSICS, UNIVERSITY OF ADELAIDE, ADELAIDE
SA 5005, AUSTRALIA

Current address: Department of physiology, University of Sydney NSW 2006, AUSTRALIA

E-mail address: nhalmagy@physiol.usyd.edu.au